

Kerr initial data

Alfonso García-Parrado Gómez-Lobo

Matematiska institutionen, Linköpings Universitet,
SE-581 83 Linköping, Sweden.

E-mail: algar@mai.liu.se

Juan A. Valiente Kroon

School of Mathematical Sciences, Queen Mary, University of London,
Mile End Road, London E1 4NS, UK.

E-mail: j.a.valiente-kroon@qmul.ac.uk

Abstract. Exploiting a 3+1 analysis of the Mars-Simon tensor, conditions on a vacuum initial data set ensuring that its development is isometric to a subset of the Kerr spacetime are found. These conditions are expressed in terms of the vanishing of a positive scalar function defined on the initial data hypersurface. Applications of this result are discussed.

PACS numbers: 04.20.Ex, 04.20-q, 04.70-s

Submitted to: *Class. Quantum Grav.*

1. Introduction

The Kerr spacetime is one of the most important exact solutions to the Einstein vacuum equations. Its relevance stems from the uniqueness theorems for black holes which state that under rather general conditions the Kerr spacetime is the only asymptotically flat, stationary, vacuum black hole solution —see e.g. the introduction of [19] for a critical review of the issue of black hole uniqueness and the involved assumptions. Although, the Kerr spacetime is very well understood from a spacetime perspective, the same cannot be said if one adopts a 3+1 point of view —which would be the case if one tries to numerically calculate the spacetime from some Cauchy initial data.

As soon as one moves away from a 3+1 gauge which is adapted to the stationary and axial symmetries —which can occur in some applications, in particular in numerical ones— an analysis of the Kerr spacetime and initial data sets thereof becomes very complicated. The explicit nature of the Kerr solution makes it tempting to perform detailed calculations in order to, say, verify a particular property of the spacetime. This approach usually leads to very lengthy expressions which can be very hard to analyse.

In a sense one could say that exact solutions contain too much information. In this case it can be more convenient to adopt more abstract approaches and, accordingly, it may prove useful to have at hand a characterization of Kerr initial data.

The question of providing an invariant characterization of initial data sets for the Schwarzschild spacetime has been addressed in [28, 16]. In particular, the analysis of [16] provides an algorithmic characterization of Schwarzschild data. That is, a procedure is provided to verify whether a given initial data set for the Einstein field equations will render a development which is isometric to a portion of the Schwarzschild spacetime.

One of the most important algebraic properties of the Kerr spacetime is that its Weyl tensor is of Petrov type D. The close relation between vacuum spacetimes with a Weyl tensor of Petrov type D and Killing spinors has been exploited in [17] to provide a characterization of initial data sets whose developments will be of Petrov type D. This characterization relies on one being able to decide whether a set of overdetermined partial differential equations has solutions for a given initial data set. Accordingly, such a characterization is not algorithmic. Although not explicitly stated in [17], from that analysis it should be possible to obtain a characterization of Kerr initial data by adding some global conditions.

The characterization of initial data sets discussed in [28, 16, 17] has followed the general strategy of starting from a given tensorial (respectively, spinorial) *spacetime* characterization of the class of spacetimes under consideration. Necessary conditions on the initial data set are obtained by performing a 3+1 decomposition of the spacetime characterization. Given a set of necessary conditions on the initial data, it is then natural to address the question of sufficiency. This is, usually, the most challenging part of the procedure as one has to discuss the evolution of complicated tensorial objects. The idea behind this is to show that if the necessary conditions are satisfied on some open subset of the initial hypersurface, then one can —possibly, under some additional assumptions— recover the spacetime characterization on the development of the open subset on the initial hypersurface from which one started.

In this article a particular characterization of Kerr initial data is addressed. Our starting point is a certain spacetime characterization of the Kerr spacetime given in [21, 22]. This characterization was developed with the aim of providing an alternative way of proving the uniqueness of Kerr spacetime among the class of stationary, asymptotically flat black holes. This expectation has been recently fulfilled in [19], where a proof of the uniqueness of Kerr which does not make assumptions on the analyticity of the metric has been given. At the heart of the spacetime characterization given in [21, 22] —cfr. theorem 3— is a certain tensor, *the Mars-Simon tensor*, whose construction requires the existence of a timelike Killing vector. The Mars-Simon tensor is a spacetime version of the *Simon tensor*, a tensor, defined in the quotient manifold of a stationary spacetime, which characterizes the Kerr spacetime —see [27].

Following the general strategy for the construction of characterizations of initial data sets out of spacetime characterizations, necessary conditions for Kerr initial data are deduced from a 3+1 splitting of the Mars-Simon tensor. Accordingly, one assumes

that the spacetime one is working with has a timelike Killing vector. This requirement can be encoded in the initial data by requiring that the data has a suitable *Killing initial data* (KID). The Mars-Simon tensor has the same symmetries as the Weyl tensor, and hence its 3+1 decomposition can be given in terms of its *electric* and *magnetic* parts. In order to discuss the propagation of the Mars-Simon tensor we make use of a framework for discussing the propagation tensorial fields using *superenergy-type* tensors —this framework has been discussed in e.g. [4]. It should be pointed out that the characterization discussed in this article is not algorithmic. That is, like the one for type D initial data discussed in [17] it depends on being able to decide whether a certain overdetermined system of partial differential equations admits a solution.

The article is structured as follows: in section 2 our main conventions are fixed and relevant aspects of the 3+1 formalism are discussed. Section 3 discusses the properties and causal propagation of Weyl candidates —i.e. tensors with the same symmetries of the Weyl tensor. Section 4 is concerned with the properties of the Mars-Simon tensor. Section 5 discusses the causal propagation of the Mars-Simon tensor. Section 6 applies the previous discussion to the construction of a characterization of Kerr initial data. Our main result is provided in theorem 6. In section 6.1 we particularize to the case of the Schwarzschild spacetime, where, with the aim of the results of [16] it is possible to obtain an initial data characterization which is algorithmic. Some concluding remarks are provided in section 7. Finally, some technical details, too lengthy to be presented in the main text are presented in the appendix A.

All the tensor computations of this paper have been performed with the software *xAct* [23]. *xAct* is a suite of MATHEMATICA packages which has among its many features the capability to efficiently canonicalize tensor expressions by the use of powerful algorithms based on permutation group theory [24]. Currently no other software, either free or commercial, is capable to handle the tensor computations needed in this paper.

2. Preliminaries

Let $(\mathcal{M}, g_{\mu\nu})$ denote a smooth orientable spacetime. The following conventions will be used: plain Greek letters $\alpha, \beta, \gamma, \dots$ denote abstract indices and boldface Latin characters $\mathbf{a}, \mathbf{b}, \mathbf{c}, \dots$ will be used for component indices. The signature of the metric tensor $g_{\mu\nu}$ will be taken to be $(-, +, +, +)$, while $R_{\mu\nu\alpha\rho}$, $R_{\mu\nu} = R^\alpha_{\mu\alpha\nu}$ and $W_{\alpha\beta\mu\nu}$ denote, respectively the Riemann, Ricci and Weyl tensors of $g_{\mu\nu}$. The tensor $\eta_{\alpha\beta\sigma\nu}$ is the volume element which is used to define the Hodge dual of any antisymmetric tensor —denoted by attaching a star $*$ to the tensor symbol. Sometimes we will need to work with complex tensors in which case the complex conjugation of a tensor is denoted by an overbar. The operator $\mathcal{L}_{\vec{u}}$ symbolizes the Lie derivative with respect to the vector field u^μ .

2.1. The orthogonal splitting

Let n^μ be a unit timelike vector, $n^\mu n_\mu = -1$ defined on $\mathcal{M}$. Then any tensor or tensorial expression can be decomposed with respect to n^μ and the way to achieve it is the essence of the orthogonal splitting (also known as 3+1 formalism) which is described in many places of the literature —see e.g. [11, 12]). We review the parts of this formalism needed in this work. The spatial metric is defined by $h_{\mu\nu} \equiv g_{\mu\nu} + n_\mu n_\nu$ and it has the algebraic properties $h^\mu{}_\mu = 3$, $h^\mu{}_\sigma h_{\sigma\nu} = h_{\mu\nu}$. We shall call a covariant tensor $T_{\alpha_1 \dots \alpha_m}$ *spatial* with respect to $h_{\mu\nu}$ if it is invariant under $h^\mu{}_\nu$ i.e. if

$$h^{\alpha_1}{}_{\beta_1} \dots h^{\alpha_m}{}_{\beta_m} T_{\alpha_1 \dots \alpha_m} = T_{\beta_1 \dots \beta_m}, \quad (2.1)$$

with the obvious generalization for any mixed tensor. This property implies that the inner contraction of n^μ with $T_{\alpha_1 \dots \alpha_m}$ (taken on any index) vanishes. The orthogonal splitting of a tensor expression consists in writing it as a sum of terms which are tensor products of the unit normal and spatial tensors of lesser degree —or the same degree in which case the unit normal is absent.

In order to find the orthogonal splitting of expressions containing covariant derivatives we need to introduce the *spatial derivative* D_μ which is an operator whose action on any tensor field $T^{\alpha_1 \dots \alpha_p}{}_{\beta_1 \dots \beta_q}$, $p, q \in \mathbb{N}$ is given by

$$D_\mu T^{\alpha_1 \dots \alpha_p}{}_{\beta_1 \dots \beta_q} \equiv h^{\alpha_1}{}_{\rho_1} \dots h^{\alpha_p}{}_{\rho_p} h^{\sigma_1}{}_{\beta_1} \dots h^{\sigma_q}{}_{\beta_q} h^\lambda{}_\mu \nabla_\lambda T^{\rho_1 \dots \rho_p}{}_{\sigma_1 \dots \sigma_q}. \quad (2.2)$$

From equation (2.2) it is clear that $D_\mu T^{\alpha_1 \dots \alpha_p}{}_{\beta_1 \dots \beta_q}$ is spatial.

The results just described hold for an arbitrary unit timelike vector n^μ but in our framework we only need to consider integrable timelike vectors which are characterized by the condition $n_{[\mu} \nabla_\nu n_{\sigma]} = 0$ (Frobenius condition). In this case there exists a foliation of $\mathcal{M}$ such that the vector field n^μ is orthogonal to the leaves of the foliation. We shall denote by $\{\Sigma_t\}$, $t \in I \subset \mathbb{R}$, the family of leaves of this foliation and Σ_0 is called the *initial data hypersurface* —it is assumed that $0 \in I$. The tensor $h_{\mu\nu}$ plays the role of the *first fundamental form* for any of the leaves while the symmetric tensor $K_{\mu\nu}$ defined by

$$K_{\mu\nu} \equiv -\frac{1}{2} \mathcal{L}_{\vec{n}} h_{\mu\nu}, \quad (2.3)$$

can be identified with the *second fundamental form*. Combining the previous definition with the Frobenius condition we easily derive

$$\nabla_\mu n_\nu = -K_{\mu\nu} - n_\mu A_\nu, \quad (2.4)$$

where $A^\mu \equiv n^\rho \nabla_\rho n^\mu$ is the acceleration of n^μ . By using these quantities one is in principle able to work out the orthogonal splitting of any tensorial expression. We supply below the explicit results of calculations which will be needed repeatedly in the sequel.

- Orthogonal splitting of the volume element:

$$\eta_{\alpha\beta\gamma\delta} = -n_\alpha \varepsilon_{\beta\gamma\delta} + n_\beta \varepsilon_{\alpha\gamma\delta} - n_\gamma \varepsilon_{\alpha\beta\delta} + n_\delta \varepsilon_{\alpha\beta\gamma}. \quad (2.5)$$

Here $\varepsilon_{\alpha\beta\gamma} \equiv n^\mu \eta_{\mu\alpha\beta\gamma}$ is the *spatial volume element* which is a fully antisymmetric spatial tensor.

- Orthogonal splitting of an antisymmetric tensor. If $H_{\mu\nu}$ is an arbitrary antisymmetric tensor then its orthogonal splitting takes the form

$$H_{\mu\nu} = 2P_{[\nu}n_{\mu]} - Q_\rho \varepsilon_{\mu\nu}{}^\rho, \quad (2.6)$$

where $P_\mu \equiv H_{\mu\nu}n^\nu$, $Q_\mu \equiv H_{\mu\nu}^*n^\nu$ are clearly spatial tensors. It is straightforward from the previous equation to find an analogous formula for the orthogonal splitting of $H_{\mu\nu}^*$. If we work with the *self-dual* part of $H_{\mu\nu}$, denoted by $\mathcal{H}_{\mu\nu}$ and defined by

$$\mathcal{H}_{\mu\nu} \equiv H_{\mu\nu} + i H_{\mu\nu}^*, \quad (2.7)$$

then its orthogonal splitting takes the form

$$\mathcal{H}_{\mu\nu} = i \varepsilon_{\rho\mu\nu} \mathcal{P}^\rho - 2\mathcal{P}_{[\mu}n_{\nu]}, \quad \mathcal{P}_\mu \equiv P_\mu + i Q_\mu. \quad (2.8)$$

- Orthogonal splitting of the Weyl tensor:

$$W_{\mu\nu\lambda\rho} = 2(l_{\mu[\lambda}E_{\rho]\nu} - l_{\nu[\lambda}E_{\rho]\mu} - n_{[\lambda}B_{\rho]\tau}\epsilon^\tau_{\mu\nu} - n_{[\mu}B_{\nu]\tau}\epsilon^\tau_{\lambda\rho}), \quad (2.9)$$

where

$$E_{\tau\rho} \equiv W_{\tau\nu\rho\lambda}n^\nu n^\lambda, \quad B_{\tau\rho} \equiv W_{\tau\nu\rho\lambda}^*n^\nu n^\lambda, \quad (2.10)$$

denote the Weyl tensor *electric* and *magnetic* parts respectively, and $l_{\mu\nu} \equiv h_{\mu\nu} + n_\mu n_\nu$. The tensors $E_{\mu\nu}$ and $B_{\mu\nu}$ are symmetric, traceless, and spatial. Equation (2.9) also holds for any rank-4 tensor possessing the same algebraic properties as the Weyl tensor. Such tensors will be designated collectively as *Weyl candidates*—see section 3. We can also choose to work with the self-dual Weyl tensor $\mathcal{W}_{\mu\nu\lambda\rho}$ which is given by

$$\mathcal{W}_{\mu\nu\lambda\rho} \equiv \frac{1}{2}(W_{\mu\nu\lambda\rho} + i W_{\mu\nu\lambda\rho}^*), \quad (2.11)$$

Using (2.9) and the analogous formula for the orthogonal splitting of $W_{\mu\nu\lambda\rho}^*$ one can obtain an expression for the orthogonal splitting of $\mathcal{W}_{\mu\nu\lambda\rho}$

$$\mathcal{W}_{\mu\nu\rho\lambda} = 2(l_{\nu[\lambda}\mathcal{E}_{\rho]\mu} + l_{\mu[\rho}\mathcal{E}_{\lambda]\nu} + i \varepsilon_{\rho\lambda\alpha}n_{[\mu}\mathcal{E}_{\nu]}^\alpha + i \varepsilon_{\mu\nu\alpha}n_{[\rho}\mathcal{E}_{\lambda]}^\alpha), \quad (2.12)$$

where

$$\mathcal{E}_{\mu\nu} \equiv \mathcal{W}_{\mu\rho\nu\lambda}n^\rho n^\lambda = \frac{1}{2}(E_{\mu\nu} - i B_{\mu\nu}), \quad \mathcal{E}_{(\mu\nu)} = \mathcal{E}_{\mu\nu}, \quad \mathcal{E}^\rho{}_\rho = 0. \quad (2.13)$$

Again a similar formula to (2.12) holds for any Weyl candidate.

- Let T^μ be any spatial vector. Then the orthogonal splitting of $\nabla_\mu T_\nu$ is

$$\nabla_\mu T_\nu = D_\mu T_\nu - n_\mu(K_{\rho\nu}T^\rho + \mathcal{L}_{\vec{n}}T_\nu) - T^\rho A_\rho n_\mu n_\nu - n_\nu K_{\rho\mu}T^\rho. \quad (2.14)$$

Note that since T_μ is spatial then $\mathcal{L}_{\vec{n}}T_\nu$ is also spatial.

2.2. The Cauchy problem

We briefly review the standard formulation of the Cauchy problem for the vacuum Einstein equations. In this formulation one considers a 3-dimensional connected Riemannian manifold (Σ, h_{ij}) —we use small plain Latin letters $i, j, k, \dots$ for the abstract

indices of tensors on this manifold—and an isometric embedding $\phi : \Sigma \longrightarrow \mathcal{M}$. The map ϕ is an isometric embedding if $\partial_i \phi^\mu \partial_j \phi^\nu g_{\mu\nu} = h_{ij}$ where $\partial_i \phi^\mu \partial_j \phi^\nu$ realises the pullback, ϕ^* , of tensor fields from $\mathcal{M}$ to Σ . The metric h_{ij} defines a unique affine connection D_i (Levi-Civita connection) by means of the standard condition

$$D_j h_{ik} = 0. \quad (2.15)$$

The Riemann tensor of D_i is r_{ijkl} and from it we define its Ricci tensor by $r_{ij} \equiv r^l{}_{ilj}$ and its scalar curvature $r \equiv r^i{}_i$ —in Σ indices are raised and lowered with h_{ij} and its inverse h^{ij} .

Theorem 1. *Let (Σ, h_{ij}) be a Riemannian manifold and suppose that there exists a symmetric tensor field K_{ij} on it which verifies the conditions (vacuum constraints)*

$$r + K^2 - K^{ij} K_{ij} = 0, \quad (2.16a)$$

$$D^j K_{ij} - D_i K = 0, \quad (2.16b)$$

where $K \equiv K^i{}_i$. Provided that h_{ij} and K_{ij} are suitably smooth there exists an isometric embedding ϕ of Σ into a globally hyperbolic, vacuum solution $(\mathcal{N}, g_{\mu\nu})$ of the Einstein field equations. The set (Σ, h_{ij}, K_{ij}) is then called a vacuum initial data set and the spacetime $(\mathcal{N}, g_{\mu\nu})$ is the data development. Furthermore the spacelike hypersurface $\phi(\Sigma)$ is a Cauchy hypersurface in $\mathcal{N}$.

A statement of this theorem containing precise regularity conditions on h_{ij} , K_{ij} is formulated in reference [20].

Under the conditions of the theorem 1, we may construct a foliation of $\mathcal{N}$ with n^μ as the timelike unit vector which is orthogonal to the leaves and, in the above notation, set $\Sigma_0 = \phi(\Sigma)$. It is then clear that $\partial_i \phi^\mu \partial_j \phi^\nu h_{\mu\nu} = h_{ij}$. Other key properties are

$$\partial_i \phi^\mu \partial_j \phi^\nu K_{\mu\nu} = K_{ij}, \quad \partial_i \phi^\mu \partial_{j_1} \phi^{\nu_1} \cdots \partial_{j_q} \phi^{\nu_q} (D_\mu T_{\nu_1 \dots \nu_q}) = D_i (\partial_{\nu_1} \phi^{\nu_1} \cdots \partial_{\nu_q} \phi^{\nu_q} T_{\nu_1 \dots \nu_q}). \quad (2.17)$$

Using these properties and the Ricci identity for n^μ one can show that

$$E_{ij} \equiv \partial_i \phi^\mu \partial_j \phi^\nu E_{\mu\nu} = r_{ij} + K K_{ij} - K_{ik} K^k{}_j, \quad (2.18a)$$

$$B_{ij} \equiv \partial_i \phi^\mu \partial_j \phi^\nu B_{\mu\nu} = \epsilon_{(i}{}^{kl} D_{|k} K_{l|j)}. \quad (2.18b)$$

Another property which is needed later on is the following one: for any spatial tensor $P_{\alpha_1 \dots \alpha_p}$, $p \in \mathbb{N}$ with respect to n^μ we have the property

$$P_{\alpha_1 \dots \alpha_p}|_{\phi(\Sigma)} = 0 \iff \phi^*(P_{\alpha_1 \dots \alpha_p}) = 0. \quad (2.19)$$

3. Weyl candidates

Fundamental for our discussion will be tensors of rank 4 having the same algebraic properties as the Weyl tensor. More precisely, one defines a *Weyl candidate* as any rank 4 tensor $C_{\alpha\beta\gamma\delta}$ fulfilling the properties

$$C_{[\alpha\beta]\gamma\delta} = C_{\alpha\beta\gamma\delta} = C_{\gamma\delta\alpha\beta}, \quad C^\alpha{}_{\alpha\gamma\delta} = 0, \quad C_{\alpha[\beta\gamma\delta]} = 0. \quad (3.1)$$

From (3.1) we easily deduce that ${}^*C_{\alpha\beta\gamma\delta} = C_{\alpha\beta\gamma\delta}^* = (1/2)\eta_{\gamma\delta\sigma\tau}C_{\alpha\beta}{}^{\sigma\tau}$ and indeed $C_{\alpha\beta\gamma\delta}^*$ is also a Weyl candidate. Given a Weyl candidate $C_{\alpha\beta\gamma\delta}$, its *Weyl current* $J_{\alpha\beta\gamma}$ is defined by

$$J_{\beta\gamma\delta} \equiv \nabla_\alpha C_{\beta\gamma\delta}^\alpha. \quad (3.2)$$

The tensor $J_{\beta\gamma\delta}$ is trace-free, antisymmetric in the last pair of indices and it has the property $J_{[\beta\gamma\delta]} = 0$. From (3.2) we deduce the following identities

$$\nabla_{[\alpha} C_{\beta\gamma]\mu\nu} = -\frac{1}{3}\eta_{\alpha\beta\gamma\sigma} J_{\mu\nu}^{*\sigma}, \quad \nabla_\beta J_{\gamma\delta}^\beta = R_{[\gamma}{}^{\sigma\rho\mu} C_{\delta]\sigma\rho\mu}, \quad (3.3)$$

where $J_{\mu\nu}^{*\sigma} \equiv \eta_{\mu\nu}{}^{\rho\lambda} J_{\rho\lambda}^\sigma/2$. Equations (3.2)-(3.3) have counterparts involving $C_{\mu\nu\rho\sigma}^*$ which we will not discuss. Also, if we combine (3.2) and (3.3) we get a wave equation for the Weyl candidate:

$$\begin{aligned} \nabla_\alpha \nabla^\alpha C_{\beta\gamma\chi\rho} = & -2R_{\gamma}{}^\alpha{}_{[\rho}{}^\delta C_{\chi]\delta\beta\alpha} - 2R_{\chi}{}^\alpha{}_{\rho}{}^\delta C_{\beta\gamma\alpha\delta} + 2R_{[\rho}{}^\alpha C_{\chi]\alpha\beta\gamma} + 2R_{\beta}{}^\alpha{}_{[\rho}{}^\delta C_{\chi]\delta\gamma\alpha} \\ & + \eta_{\chi\rho\lambda\pi} \nabla^\pi J_{\beta\gamma}^{*\lambda} + 2\nabla_{[\chi} J_{\rho]\beta\gamma}. \end{aligned} \quad (3.4)$$

The previous considerations hold regardless of whether the Weyl candidate is real or *complex*. Indeed, from any real Weyl candidate $C_{\mu\nu\rho\sigma}$ we may construct a complex Weyl candidate $\mathcal{C}_{\mu\nu\rho\sigma}$ by means of

$$\mathcal{C}_{\mu\nu\rho\sigma} = \frac{1}{2}(C_{\mu\nu\rho\sigma} + i C_{\mu\nu\rho\sigma}^*). \quad (3.5)$$

A Weyl candidate constructed in such a way is *self-dual*[‡]

$$\mathcal{C}_{\mu\nu\rho\sigma}^* = i \mathcal{C}_{\mu\nu\rho\sigma}. \quad (3.6)$$

Reciprocally, if a complex Weyl candidate $\mathcal{C}_{\mu\nu\rho\sigma}$ fulfils (3.6) then its real part $\text{Re}(\mathcal{C}_{\mu\nu\rho\lambda})$ and its imaginary part $\text{Im}(\mathcal{C}_{\mu\nu\rho\lambda})$ are also Weyl candidates related via

$$\text{Im}(\mathcal{C}_{\mu\nu\rho\lambda}) = \frac{1}{2}\eta_{\rho\lambda\alpha\beta}\text{Re}(\mathcal{C}_{\mu\nu}{}^{\alpha\beta}), \quad \text{Re}(\mathcal{C}_{\mu\nu\rho\lambda}) = -\frac{1}{2}\eta_{\rho\lambda\alpha\beta}\text{Im}(\mathcal{C}_{\mu\nu}{}^{\alpha\beta}). \quad (3.7)$$

3.1. The causal propagation of a Weyl candidate

Besides the basic algebraic properties of a Weyl candidate just explained we need to introduce a further concept which will play a crucial role in the sequel.

Definition 1. Let $(\mathcal{M}, g_{\mu\nu})$ be a spacetime and consider a Weyl candidate $C_{\alpha\beta\mu\nu}$ defined in the whole of $\mathcal{M}$. We say that the Weyl candidate $C_{\alpha\beta\mu\nu}$ propagates causally on $\mathcal{M}$ if for any embedded spacelike hypersurface $\mathcal{B} \subset \mathcal{M}$ the condition $C_{\alpha\beta\mu\nu}|_{\mathcal{B}} = 0$ implies $C_{\alpha\beta\mu\nu} = 0$ on $D(\mathcal{B})$.

Given a set $\mathcal{B}$, a point $p \in \mathcal{M}$ belongs to the future Cauchy development $D^+(\mathcal{B})$ if any past-inextendible causal curve containing p intersects $\mathcal{B}$. There is a similar notion of $D^-(\mathcal{B})$ and the set $D(\mathcal{B}) \equiv D^+(\mathcal{B}) \cup D^-(\mathcal{B})$ is called the *Cauchy development* of $\mathcal{B}$. It is well-known that the interior of $D(\mathcal{B})$ is globally hyperbolic and since $\mathcal{B}$ is an embedded

[‡] Note that sometimes the opposite convention is followed in the literature whereby the complex conjugate of $\mathcal{C}_{\mu\nu\rho\sigma}$ is given the name of “self-dual”.

spacelike hypersurface then it will be a Cauchy hypersurface for $D(\mathcal{B})$ —see e.g. [29] for an elementary introduction to these concepts of causality theory.

The simplest example of a Weyl candidate which propagates causally is the Weyl tensor $W_{\alpha\beta\mu\nu}$ of a vacuum spacetime —see e.g. [5]. It is possible to address the causal propagation of a Weyl candidate by following the general techniques given in [4] —see also [26]— in which the causal propagation of any tensor is analysed. These ideas are reviewed next.

An essential object to study the causal propagation of a Weyl candidate is its *Bel-Robinson tensor*. Given a real Weyl candidate $C_{\alpha\beta\gamma\delta}$, we define its Bel-Robinson tensor by

$$T_{\alpha\beta\gamma\delta} \equiv C_{\alpha}{}^{\rho}{}_{\delta}{}^{\mu} C_{\beta\rho\gamma\mu} + C_{\alpha}{}^{\mu}{}_{\gamma}{}^{\rho} C_{\beta\mu\delta\rho} - \frac{1}{8} g_{\alpha\beta} g_{\gamma\delta} C_{\mu\nu\rho\lambda} C^{\mu\nu\rho\lambda}. \quad (3.8)$$

The previous definition has been described in a more general framework in [25] as the *superenergy tensor* of the Weyl candidate $C_{\alpha\beta\gamma\delta}$. In our particular context we prefer to call this tensor the Bel-Robinson tensor of $C_{\alpha\beta\gamma\delta}$ in analogy with the Bel-Robinson tensor constructed out of the Weyl tensor [3]. The Bel-Robinson tensor of a Weyl candidate has the following properties —see [25] for detailed proofs of points (i), (ii) and (iii); point (iv) follows by taking the covariant divergence of (3.8) and then using repeatedly (3.2) and (3.3).

Theorem 2. *If $T_{\alpha\beta\mu\nu}$ is the Bel-Robinson tensor of the Weyl candidate $C_{\alpha\beta\mu\nu}$ then*

- (i) $T_{(\alpha\beta\mu\nu)} = T_{\alpha\beta\mu\nu}$, $T^{\alpha}{}_{\alpha\mu\nu} = 0$.
- (ii) *Generalized dominant property: if u_1^{μ} , u_2^{μ} , u_3^{μ} , u_4^{μ} are causal future-directed vectors then $T_{\alpha\beta\mu\nu} u_1^{\alpha} u_2^{\beta} u_3^{\mu} u_4^{\nu} \geq 0$. This property admits an alternative formulation: let $E = \{e_0^{\mu}, e_1^{\mu}, e_2^{\mu}, e_3^{\mu}\}$ be any orthonormal frame with e_0^{μ} the timelike vector. Then the following inequality holds*

$$T_{0000} \geq |T_{abcd}|,$$

where the component indices refer to the frame E .

- (iii) $T_{\alpha\beta\mu\nu} = 0 \iff C_{\alpha\beta\mu\nu} = 0 \iff \exists$ a timelike vector u^{μ} such that $T_{\alpha\beta\mu\nu} u^{\alpha} u^{\beta} u^{\mu} u^{\nu} = 0$.
- (iv) The covariant divergence of $T_{\alpha\beta\mu\nu}$ is given by

$$\nabla_{\alpha} T^{\alpha}{}_{\beta\gamma\delta} = 4 J_{(\gamma}{}^{\sigma} C_{\beta)\alpha\delta\sigma} - 2 J_{(\gamma}{}^{\alpha\sigma} C_{\beta)\delta\alpha\sigma} - g_{\beta\gamma} J^{\alpha\rho\sigma} C_{\delta\alpha\sigma\rho}. \quad (3.9)$$

Next, we explain how the properties summarized in theorem 2 are useful in the study of the causal propagation of $C_{\mu\nu\rho\lambda}$. We follow closely [4] in our exposition. Let $D^+(\mathcal{B})$ be the future Cauchy development of a closed achronal hypersurface $\mathcal{B}$ and pick up any event $q \in D^+(\mathcal{B})$. Furthermore, let $J^-(q)$ denote the causal past of the event q . The set $\mathcal{K} = J^-(q) \cap \overline{D^+(\mathcal{B})}$ is compact and it contains the set $\tilde{\mathcal{B}} \equiv \mathcal{K} \cap \mathcal{B}$. The boundary of $\mathcal{K}$ is $\partial\mathcal{K} = \tilde{\mathcal{B}} \cup H^+(\tilde{\mathcal{B}})$ where $H^+(\tilde{\mathcal{B}})$ is the future Cauchy horizon of $\tilde{\mathcal{B}}$. Also, since the interior of $D^+(\mathcal{B})$ is globally hyperbolic we deduce that it can be foliated by a family of spacelike hypersurfaces $\{\Sigma_t\}$, $t \in I$, where $t_0 \leq t \leq t_1$, $t_0 < 0$, $0 < t_1$ and $\Sigma_0 = \mathcal{B}$. The figure 3.1 shows a schematic view of this geometric construction.




Figure 1. Schematic depiction of the sets $\mathcal{B}$, $\tilde{\mathcal{B}}$, $D^+(\mathcal{B})$ and $H^+(\tilde{\mathcal{B}})$ —see the main text for further details.

Let n_μ be the timelike normal 1-form to the leaves of $\{\Sigma_t\}_{t \in I}$ and define the quantity

$$W(t) \equiv \int_{J^-(\Sigma_t) \cap \mathcal{K}} T_{\mu\nu\alpha\beta} n^\mu n^\nu n^\alpha n^\beta d\mathcal{M} = \int_0^t dt' \left(\int_{\Sigma_{t'} \cap \mathcal{K}} T_{\mu\nu\alpha\beta} n^\mu n^\nu n^\alpha n^\beta d\Sigma_{t'} \right). \quad (3.10)$$

Here, $d\mathcal{M}$ is the positive measure on the spacetime $\mathcal{M}$ constructed from the volume form $\eta_{\mu\nu\rho\sigma}$ (any measure defined from a volume form will be understood as positive) and $d\Sigma_{t'}$ is the measure on the hypersurface $\Sigma_{t'}$ obtained from its volume form $n^\mu \eta_{\mu\nu\rho\sigma}|_{\Sigma_{t'}}$. The scalar $T_{\mu\nu\rho\lambda} n^\mu n^\nu n^\rho n^\lambda$ is called the *superenergy density* and is everywhere non-negative —see point (ii) of theorem 2— which entails $W(t) \geq 0$. Also, from the definition of $W(t)$ we get

$$W'(t) = \int_{\Sigma_t \cap \mathcal{K}} T_{\mu\nu\alpha\beta} n^\mu n^\nu n^\alpha n^\beta d\Sigma_t = \int_{\Sigma_t \cap \mathcal{K}} P_\mu n^\mu d\Sigma_t, \quad (3.11)$$

where P_μ is defined by

$$P_\mu \equiv T_{\mu\nu\rho\sigma} n^\nu n^\rho n^\sigma. \quad (3.12)$$

Again point (ii) of theorem 2 implies that $P_\mu n^\mu$ is also a non-negative quantity and hence $W'(t) \geq 0$. Now, according to the Gauss theorem we have

$$\int_{J^-(\Sigma_t) \cap \mathcal{K}} \nabla_\mu P^\mu d\mathcal{M} = \int_{\Sigma_t \cap \mathcal{K}} P_\mu n^\mu d\Sigma_t - \int_{\tilde{\mathcal{B}}} P_\mu n^\mu d\tilde{\mathcal{B}} + \int_{H^+(\tilde{\mathcal{B}})} P_\mu k^\mu d(H^+(\tilde{\mathcal{B}})), \quad (3.13)$$

where $d\tilde{\mathcal{B}}$ and $d(H^+(\tilde{\mathcal{B}}))$ denote the measures on the hypersurfaces $\tilde{\mathcal{B}}$ and $H^+(\tilde{\mathcal{B}})$ respectively induced by the volume forms $n^\mu \eta_{\mu\nu\rho\lambda}|_{\tilde{\mathcal{B}}}$, and $k^\mu \eta_{\mu\nu\rho\lambda}|_{H^+(\tilde{\mathcal{B}})}$. In this last expression k^μ is a causal vector which is orthogonal to the null hypersurface $H^+(\tilde{\mathcal{B}})$, points outward to $\mathcal{K}$ (therefore it is future-directed) and is such that $k^\mu \eta_{\mu\nu\rho\lambda}|_{H^+(\tilde{\mathcal{B}})} \neq 0$. Note that these three conditions do not fix the vector k^μ univocally. By combining (3.11) and (3.13) we deduce

$$W'(t) = \int_{J^-(\Sigma_t) \cap \mathcal{K}} \nabla_\mu P^\mu d\mathcal{M} + \int_{\tilde{\mathcal{B}}} P^\mu n_\mu d\tilde{\mathcal{B}} - \int_{H^+(\tilde{\mathcal{B}}) \cap J^-(\Sigma_t)} P^\mu k_\mu d(H^+(\tilde{\mathcal{B}})). \quad (3.14)$$

Now, we need to estimate the right hand side of (3.14). According to point (ii) of theorem 2, the quantity $P^\mu k_\mu$ is positive and therefore

$$0 \leq \int_{H^+(\tilde{\mathcal{B}}) \cap J^-(\Sigma_t)} P^\mu k_\mu d(H^+(\tilde{\mathcal{B}})). \quad (3.15)$$

On the other hand, if $C_{\alpha\beta\mu\nu}$ vanishes on $\mathcal{B}$ then $W(0) = 0$ and

$$\int_{\tilde{\mathcal{B}}} P^\mu n_\mu d\tilde{\mathcal{B}} = 0. \quad (3.16)$$

Next, we set up an orthonormal frame $E = \{e_0^\mu, e_1^\mu, e_2^\mu, e_3^\mu\}$ with $e_0^\mu = n^\mu$ and we consider the set of components $\nabla_a n_b$ in such a frame. Since these are continuous functions we deduce that in the compact set $\mathcal{K}$ the estimate $\nabla_a n_b \leq m_0$, $\forall a, b = 0, 1, 2, 3$ holds for some constant m_0 . Hence

$$\begin{aligned} \nabla_a P^a &= 3n_c n_d T^{cdab} \nabla_a n_b + n_a n_b n_c \nabla_d T^{dabc} \\ &\leq 3m_0 \sum_{c,d=0}^3 T^{00cd} + n_a n_b n_c \nabla_d T^{dabc} \leq 3m_0 T^{abcd} n_a n_b n_c n_d + n_a n_b n_c \nabla_d T^{dabc}, \end{aligned} \quad (3.17)$$

where in the last step, point (ii) of theorem 2 was used. From here, we deduce

$$\begin{aligned} &\int_{J^-(\Sigma_t) \cap \mathcal{K}} \nabla_\mu P^\mu d\mathcal{M} \\ &\leq \int_{J^-(\Sigma_t) \cap \mathcal{K}} (\nabla_\mu T^{\mu\nu\rho\sigma}) n_\nu n_\rho n_\sigma d\mathcal{M} + 3m_0 \int_{J^-(\Sigma_t) \cap \mathcal{K}} T^{\mu\nu\rho\sigma} n_\mu n_\nu n_\rho n_\sigma d\mathcal{M}. \end{aligned} \quad (3.18)$$

Using the above estimates in (3.14) we get

$$0 \leq W'(t) \leq \int_{J^-(\Sigma_t) \cap \mathcal{K}} (\nabla_\mu T^{\mu\nu\rho\sigma}) n_\nu n_\rho n_\sigma d\mathcal{M} + 3m_0 W(t). \quad (3.19)$$

Now it only remains to estimate $n_\nu n_\rho n_\sigma \nabla_\mu T^{\mu\nu\rho\sigma}$ and to that end (3.9) is used. The actual estimate will depend on the particular expression of the Weyl current. For example, the simplest case arises when $J_{\mu\nu\rho} = 0$ where no further estimates are needed and Grönwall's lemma entails

$$W(t) = 0, \quad t \in [0, t_1], \quad (3.20)$$

which in turn implies that the scalar $T_{\mu\nu\alpha\beta} n^\mu n^\nu n^\alpha n^\beta$ vanishes in $\mathcal{K}$. From the arbitrariness of q we deduce that the superenergy density is zero in $D^+(\mathcal{B})$ and by a similar argument as the above one can show that this is also the case for $D^-(\mathcal{B})$. Hence point 3 of theorem 2 implies that $C_{\alpha\beta\mu\nu} = 0$ on $D^\pm(\mathcal{B})$. Consequently the tensor $C_{\alpha\beta\mu\nu}$ propagates causally whenever the Weyl current is zero.

Remark 1. Point (iii) of theorem 2 entails $C_{\alpha\beta\mu\nu}|_{\mathcal{B}} = 0$ if and only if $(T_{\alpha\beta\mu\nu} n^\alpha n^\beta n^\mu n^\nu)|_{\mathcal{B}} = 0$. Accordingly, the causal propagation of a Weyl candidate can also be formulated in terms of the vanishing of the superenergy density at the hypersurface $\mathcal{B}$.

4. The Mars-Simon tensor

We start this section by recalling some basic background about vacuum spacetimes possessing a Killing vector —see e.g. [18] for a complete account. Let $(\mathcal{M}, g_{\mu\nu})$ be a vacuum solution of Einstein field equations and let ξ^μ be a Killing vector, $\nabla_\mu \xi_\nu + \nabla_\nu \xi_\mu = 0$. This condition enables us to define a 2-form, the *Killing form*, by

$$F_{\mu\nu} \equiv \nabla_{[\mu} \xi_{\nu]} = \nabla_\mu \xi_\nu. \quad (4.1)$$

Elementary manipulations lead to

$$\nabla_\mu F_{\nu\lambda} = W_{\nu\lambda\mu\rho} \xi^\rho. \quad (4.2)$$

The *self-dual Killing form* $\mathcal{F}_{\mu\nu}$ is then

$$\mathcal{F}_{\mu\nu} = F_{\mu\nu} + iF_{\mu\nu}^*, \quad (4.3)$$

with $F_{\mu\nu}^* = (1/2)\eta_{\mu\nu\lambda\rho}F^{\lambda\rho}$ the Hodge dual of $F_{\mu\nu}$. From the previous definition it is straightforward to obtain the property (self-duality)

$$\mathcal{F}_{\mu\nu} = i \mathcal{F}_{\mu\nu}^*. \quad (4.4)$$

We shall write $\mathcal{F}^2 = \mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu}$. The Killing form is said to be degenerate if $\mathcal{F}^2 = 0$. Other basic algebraic properties of $\mathcal{F}_{\mu\nu}$ are

$$\mathcal{F}_{\mu\rho}\mathcal{F}_\nu{}^\rho = \frac{1}{4}g_{\mu\nu}\mathcal{F}^2, \quad \mathcal{F}_{[\mu}{}^\rho\overline{\mathcal{F}}_{\nu]\rho} = 0, \quad \mathcal{F}_{\mu\nu}\overline{\mathcal{F}}^{\mu\nu} = 0. \quad (4.5)$$

From a Killing vector ξ^μ we may define its associated *Ernst 1-form* by

$$\sigma_\nu \equiv 2\xi^\mu \mathcal{F}_{\mu\nu} = -\nabla_\nu \Lambda - i\Omega_\nu, \quad (4.6)$$

where $\Lambda \equiv \xi_\mu \xi^\mu$ is the norm of the Killing vector and $\Omega_\mu \equiv \eta_{\mu\nu\lambda\rho} \xi^\nu \nabla^\lambda \xi^\rho$ denotes its *twist form*. It is well-known that in a vacuum spacetime σ_μ is closed —i.e. $\nabla_{[\nu} \sigma_{\mu]} = 0$. Thus locally —if we are in a simply connected region— there exists a scalar field σ , the *Ernst potential*, such that $\sigma_\mu = \nabla_\mu \sigma$. The self-dual Weyl tensor and the self-dual Killing form are related by

$$\nabla_\mu \mathcal{F}_{\nu\rho} = 2\overline{\mathcal{W}}_{\nu\rho\mu\alpha} \xi^\alpha, \quad (4.7)$$

a property which is easily obtained from (4.2). Finally, we introduce the tensor $\mathcal{I}_{\mu\nu\lambda\rho}$ defined by

$$\mathcal{I}_{\mu\nu\lambda\rho} \equiv \frac{1}{2}(g_{\mu\lambda}g_{\nu\rho} - g_{\mu\rho}g_{\nu\lambda} + i\eta_{\mu\nu\lambda\rho}). \quad (4.8)$$

The tensor $\mathcal{I}_{\mu\nu\lambda\rho}$ has the obvious symmetries $\mathcal{I}_{\mu\nu\lambda\rho} = \mathcal{I}_{[\mu\nu]\lambda\rho} = \mathcal{I}_{\lambda\rho\mu\nu}$. This tensor can be regarded as a metric in the space of self-dual 2-forms.

We have now all the ingredients to define the *Mars-Simon tensor*. This is a four-rank tensor $\mathcal{S}_{\mu\nu\lambda\rho}$ whose expression is given by §

$$\mathcal{S}_{\mu\nu\lambda\rho} \equiv 2\overline{\mathcal{W}}_{\mu\nu\lambda\rho} + \frac{6}{1-\sigma} \left(\mathcal{F}_{\mu\nu}\mathcal{F}_{\lambda\rho} - \frac{1}{6}\mathcal{F}^2\mathcal{I}_{\mu\nu\lambda\rho} \right). \quad (4.9)$$

§ We follow the convention set in [19] of calling this tensor the Mars-Simon tensor. Note however, that our conventions are slightly different.

From the previous expression we deduce that in a vacuum spacetime admitting a Killing vector, the Mars-Simon tensor is determined up to a choice for the Ernst potential. We make a choice such that $\sigma \neq 1$ on $\mathcal{M}$. Our interest in the Mars-Simon tensor lies in the following result —see [21, 22] for a proof.

Theorem 3. *Let $(\mathcal{M}, g_{\mu\nu})$ be a smooth, vacuum spacetime with a Killing vector ξ^μ . Let $\mathcal{F}_{\mu\nu}$ denote the associated self-dual Killing form. If there is a non-vanishing real constant M such that the conditions*

$$\mathcal{F}^2 = -M^2(1 - \sigma)^4, \quad (4.10a)$$

$$\mathcal{S}_{\mu\nu\lambda\rho} = 0, \quad (4.10b)$$

hold on a non-empty $\mathcal{N} \subset \mathcal{M}$ then $(\mathcal{N}, g_{\mu\nu})$ is locally isometric to the Kerr spacetime.

Condition (4.10a) of theorem 3 can be reformulated in the form

$$\Xi(M, \vec{\xi}) = 0, \quad (4.11)$$

where the scalar $\Xi(M, \vec{\xi})$ is defined for any Killing vector $\vec{\xi}$ and any real constant M by the expression

$$\Xi(M, \vec{\xi}) \equiv \mathcal{F}^2 + M^2(1 - \sigma)^4. \quad (4.12)$$

A very important property of the scalar $\Xi(M, \vec{\xi})$ is that it fulfils the differential equation

$$\nabla_\mu \Xi = -2\mathcal{F}^{\alpha\beta} \mathcal{S}_{\mu\rho\alpha\beta} \xi^\rho + \frac{4\Xi\sigma_\mu}{\sigma - 1}. \quad (4.13)$$

To see it, differentiate both sides of (4.12) and use (4.7) to replace the covariant derivatives of $\mathcal{F}_{\mu\nu}$. The result is

$$\nabla_\mu \Xi = -4(\mathcal{F}^{\alpha\beta} \overline{\mathcal{W}}_{\mu\rho\alpha\beta} \xi^\rho + M^2 \sigma_\mu (1 - \sigma)^3). \quad (4.14)$$

Next use (4.9) to write $\overline{\mathcal{W}}_{\mu\rho\alpha\beta}$ in terms of $\mathcal{S}_{\mu\rho\alpha\beta}$ and replace all the occurrences of $\mathcal{I}_{\mu\rho\alpha\beta}$ by means of (4.8) to obtain

$$\nabla_\mu \Xi = 2(2M^2 \sigma_\mu (\sigma - 1)^3 - \mathcal{F}^{\alpha\beta} \mathcal{S}_{\mu\rho\alpha\beta} \xi^\rho) + \frac{\mathcal{F}^2(-10\mathcal{F}_{\mu\rho} + 2i \mathcal{F}_{\mu\rho}^*) \xi^\rho}{\sigma - 1}. \quad (4.15)$$

We use here (4.4) and (4.6) with the result

$$\nabla_\mu \Xi = -2\mathcal{F}^{\alpha\beta} \mathcal{S}_{\mu\rho\alpha\beta} \xi^\rho + \frac{4\sigma_\mu (\mathcal{F}^2 + M^2(1 - \sigma)^4)}{\sigma - 1}, \quad (4.16)$$

from which (4.13) is deduced.

4.1. Mars-Simon tensor as a Weyl candidate

In [19] it is proved that the Mars-Simon tensor $\mathcal{S}_{\mu\nu\alpha\beta}$ is a Weyl candidate. Furthermore, it was shown that its Weyl current is a linear function of $\mathcal{S}_{\mu\nu\alpha\beta}$ in which case the wave equation (3.4) is homogeneous in $\mathcal{S}_{\mu\nu\alpha\beta}$. The derivation of these results are discussed as some of the intermediate calculations will be needed in section 5.

Proposition 1. *The Mars-Simon tensor is a Weyl candidate.*

Proof. From (4.9) it is clear that $\mathcal{S}_{[\mu\nu]\alpha\lambda} = \mathcal{S}_{\mu\nu[\alpha\lambda]} = \mathcal{S}_{\alpha\lambda\mu\nu}$. Using (4.5) one can show that $\mathcal{S}_{\mu\rho}{}^\rho{}_\lambda = 0$ and therefore the Mars-Simon tensor is traceless. To finish our proof we need to show that the Mars-Simon tensor fulfils the cyclic identity. To that end we compute $\mathcal{S}_{\mu[\nu\rho\lambda]}$ getting

$$\mathcal{S}_{\mu[\nu\rho\lambda]} = \frac{6}{1-\sigma}\mathcal{F}_{\mu[\nu}\mathcal{F}_{\rho\lambda]} - \frac{i}{2(1-\sigma)}\eta_{\mu\nu\rho\lambda}. \quad (4.17)$$

The right hand side of this expression vanishes by virtue of the identity

$$\mathcal{F}_{\mu[\nu}\mathcal{F}_{\rho\lambda]} = \frac{1}{12}i\mathcal{F}^2\eta_{\mu\nu\rho\lambda}, \quad (4.18)$$

which in turn is a consequence of the algebraic property $\mathcal{F}_{\mu[\nu}\mathcal{F}_{\rho\lambda]} = \mathcal{F}_{[\mu\nu}\mathcal{F}_{\rho\lambda]}$ and the fact that in four dimensions $\mathcal{F}_{[\mu\nu}\mathcal{F}_{\rho\lambda]}$ is proportional to $\eta_{\mu\nu\rho\lambda}$. $\square$

A direct computation shows that the Mars-Simon tensor is self-dual

$$\mathcal{S}_{\mu\nu\rho\lambda} = i\mathcal{S}_{\mu\nu\rho\lambda}^*. \quad (4.19)$$

Therefore $\text{Re}(\mathcal{S}_{\mu\nu\rho\lambda})$ and $\text{Im}(\mathcal{S}_{\mu\nu\rho\lambda})$ are then related by duality as explained in section 3:

$$\text{Im}(\mathcal{S}_{\mu\nu\rho\lambda}) = \frac{1}{2}\eta_{\rho\lambda\alpha\beta}\text{Re}(\mathcal{S}_{\mu\nu}{}^{\alpha\beta}), \quad \text{Re}(\mathcal{S}_{\mu\nu\rho\lambda}) = -\frac{1}{2}\eta_{\rho\lambda\alpha\beta}\text{Im}(\mathcal{S}_{\mu\nu}{}^{\alpha\beta}). \quad (4.20)$$

Any property of $\mathcal{S}_{\mu\nu\rho\lambda}$ will admit an equivalent counterpart formulated in terms of either its real or imaginary part. In particular, this shows that both $\text{Re}(\mathcal{S}_{\mu\nu\rho\lambda})$ and $\text{Im}(\mathcal{S}_{\mu\nu\rho\lambda})$ are Weyl candidates and any of them contains the same information as the original Mars-Simon tensor.

Proposition 2. *The covariant divergence of the Mars-Simon tensor is given by*

$$\nabla_\alpha \mathcal{S}^\alpha{}_{\beta\gamma\delta} = \frac{1}{\sigma-1} \left(4(\mathcal{F}_{[\delta}{}^\rho \mathcal{S}_{\gamma]\beta\alpha\rho} + \mathcal{F}_\beta{}^\rho \mathcal{S}_{\gamma\delta\alpha\rho}) + 2g_{\beta[\gamma} \mathcal{S}_{\delta]\alpha\lambda\rho} \mathcal{F}^{\lambda\rho} \right) \xi^\alpha. \quad (4.21)$$

Proof. To calculate $\nabla_\alpha \mathcal{S}^\alpha{}_{\beta\gamma\delta}$ we start from (4.9) and take the covariant divergence of $\mathcal{S}_{\mu\nu\lambda\rho}$. Next we apply the relations (4.7), $\nabla_\mu \sigma = \sigma_\mu$, $\nabla_\alpha \bar{\mathcal{W}}^\alpha{}_{\sigma\mu\nu} = 0$ and $\nabla_\alpha \mathcal{I}_{\mu\nu\rho\sigma} = 0$. Finally we expand all the occurrences of $\mathcal{I}_{\mu\nu\rho\sigma}$ using (4.8) with the result

$$\begin{aligned} \nabla_\alpha \mathcal{S}^\alpha{}_{\beta\gamma\delta} &= \frac{1}{2(1-\sigma)^2} (-12\mathcal{F}_{\beta\alpha} \mathcal{F}_{\gamma\delta} \sigma^\alpha + (i\eta_{\beta\gamma\delta\alpha} \sigma^\alpha + 2g_{\beta[\gamma} \sigma_{\delta]}) \mathcal{F}^2) \\ &+ \frac{1}{1-\sigma} ((2i\eta_{\beta\gamma\delta\alpha} \bar{\mathcal{W}}_\rho{}^\alpha{}_{\pi\kappa} \xi^\rho + 4g_{\beta[\delta} \bar{\mathcal{W}}_{\gamma]\alpha\pi\kappa} \xi^\alpha) \mathcal{F}^{\pi\kappa} - 12\mathcal{F}_\beta{}^\rho \bar{\mathcal{W}}_{\gamma\delta\alpha\rho} \xi^\alpha) \end{aligned} \quad (4.22)$$

The dependence on the Ernst 1-form can be removed from this expression if we write σ_μ in terms of $\mathcal{F}_{\mu\nu}$ and ξ^μ by means of (4.6). The final step is to rearrange the terms in the resulting expression in order to render it in the form of (4.21). This is accomplished by using (4.9) to write $\bar{\mathcal{W}}_{\mu\nu\lambda\rho}$ in terms of the Mars-Simon tensor, expanding again all the occurrences of $\mathcal{I}_{\mu\nu\rho\sigma}$ by means of (4.8) and using (4.5), (4.4) where necessary. $\square$

Equation (4.21) enables us to define the Weyl current of the Mars-Simon tensor as

$$\mathcal{J}_{\beta\gamma\delta} \equiv \frac{1}{\sigma-1} \left(4(\mathcal{F}_{[\delta}{}^\rho \mathcal{S}_{\gamma]\beta\alpha\rho} + \mathcal{F}_\beta{}^\rho \mathcal{S}_{\gamma\delta\alpha\rho}) + 2g_{\beta[\gamma} \mathcal{S}_{\delta]\alpha\sigma\rho} \mathcal{F}^{\sigma\rho} \right) \xi^\alpha. \quad (4.23)$$

Since $\mathcal{J}_{\beta\gamma\delta}$ is linear in the Mars-Simon tensor, we conclude—in view of (3.4)—that $\mathcal{S}_{\mu\nu\rho\sigma}$ fulfils a homogeneous wave equation. This interesting property was used in [19] to show that the *domain of outer communication* of an asymptotically flat smooth stationary spacetime is, under certain additional conditions, isometric to the Kerr domain of outer communication—i.e. the *uniqueness of Kerr*.

Using $\nabla_\alpha \mathcal{S}^\alpha_{\mu\nu\rho} = \mathcal{J}_{\mu\nu\rho}$ together with the self-duality of the Mars-Simon tensor one deduces

$$\mathcal{J}_{\mu\nu\rho} = \mathrm{i} \mathcal{J}_{\mu\nu\rho}^* = \frac{\mathrm{i}}{2} \eta_{\nu\rho\alpha\beta} \mathcal{J}_\mu^{\alpha\beta}, \quad (4.24)$$

that is to say, $\mathcal{J}_{\mu\nu\rho}$ is also self-dual with respect to the block of indices $\nu\rho$.

4.2. Orthogonal splitting of the Mars-Simon tensor

In our forthcoming calculations it is necessary to use the orthogonal splitting of $\mathcal{S}_{\mu\nu\rho\lambda}$ with respect to a unit timelike vector field n^μ . Since $\mathcal{S}_{\mu\nu\rho\lambda}$ is a Weyl candidate we can use (2.12) to find such a orthogonal splitting. The resulting split is given by

$$\mathcal{S}_{\mu\nu\rho\lambda} = 2(l_{\nu[\lambda} \mathcal{T}_{\rho]\mu} + l_{\mu[\rho} \mathcal{T}_{\lambda]\nu} + \mathrm{i} \varepsilon_{\rho\lambda\alpha} n_{[\mu} \mathcal{T}_{\nu]}^\alpha + \mathrm{i} \varepsilon_{\mu\nu\alpha} n_{[\rho} \mathcal{T}_{\lambda]}^\alpha), \quad (4.25)$$

where

$$\mathcal{T}_{\mu\nu} \equiv \mathcal{S}_{\mu\rho\nu\lambda} n^\rho n^\lambda. \quad (4.26)$$

The tensor $\mathcal{T}_{\mu\nu}$ is symmetric, spatial and traceless. Alternatively, the orthogonal splitting of the Mars-Simon tensor can be calculated directly from (4.9). To that end one needs to find the orthogonal splitting of the different quantities which appear in (4.9). The orthogonal splitting of $\mathcal{W}_{\mu\nu\rho\lambda}$ is given in (2.12) and the orthogonal splitting of $\mathcal{F}_{\mu\nu}$ is calculated by means of (2.8) and is given by

$$\mathcal{F}_{\mu\nu} = \mathrm{i} \varepsilon_{\rho\mu\nu} \mathcal{E}^\rho - 2\mathcal{E}_{[\mu} n_{\nu]}, \quad (4.27)$$

where

$$\mathcal{E}_\mu \equiv \mathcal{F}_{\mu\rho} n^\rho. \quad (4.28)$$

One also needs the orthogonal splitting of $\mathcal{I}_{\mu\nu\rho\lambda}$ which is easily calculated from (4.8) and (2.5) so that

$$\mathcal{I}_{\mu\nu\rho\lambda} = h_{\mu[\rho} h_{\lambda]\nu} + 2n_{[\nu} h_{\mu][\lambda} n_{\rho]} + \mathrm{i} (n_{[\mu} \varepsilon_{\nu]\rho\lambda} + n_{[\rho} \varepsilon_{\lambda]\mu\nu}). \quad (4.29)$$

We insert relations (2.12), (4.27) and (4.29) into the formula (4.9) and then equal the resulting expression to (4.25). The result of that is the relation

$$\mathcal{T}_{\mu\nu} = 2\bar{\mathcal{E}}_{\mu\nu} + \frac{2}{1-\sigma} (3\mathcal{E}_\mu \mathcal{E}_\nu - h_{\mu\nu} \mathcal{E}_\alpha \mathcal{E}^\alpha), \quad (4.30)$$

which shall be used later on. Also needed later on is the relation

$$\mathcal{F}^2 = -4\mathcal{E}_\mu \mathcal{E}^\mu \implies \Xi(M, \vec{\xi}) = -4\mathcal{E}_\mu \mathcal{E}^\mu + M^2(1-\sigma)^4, \quad (4.31)$$

which is derived from equation (4.27).

The Killing vector ξ^μ can be decomposed in the form $\xi^\mu = -Yn^\mu + Y^\mu$ where Y , Y^μ are called the *Killing lapse* and the *Killing shift* respectively. We need to find a formula relating $\mathcal{E}_\mu$ to Y , Y_μ . To that end, we start by taking the covariant derivative of both sides of $\xi_\mu = -Yn_\mu + Y_\mu$ and replace $\nabla_\mu n_\nu$ by means of (2.4). The result is

$$\nabla_\mu \xi_\nu = Y(K_{\mu\nu} + A_\nu n_\mu) + \nabla_\mu Y_\nu - n_\nu \nabla_\mu Y.$$

Next we use in this expression the property $\nabla_\mu Y = D_\mu Y - n_\mu \mathcal{L}_{\vec{n}} Y$ and equation (2.14) to replace $\nabla_\mu Y$ and $\nabla_\mu Y_\nu$ respectively yielding

$$\begin{aligned} \nabla_\mu \xi_\nu &= Y K_{\mu\nu} + D_\mu Y_\nu + n_\mu n_\nu (\mathcal{L}_{\vec{n}} Y - A^\rho Y_\rho) \\ &\quad + n_\mu (Y A_\nu - K_{\nu\rho} Y^\rho - \mathcal{L}_{\vec{n}} Y_\nu) - n_\nu (K_{\mu\rho} Y^\rho + D_\mu Y). \end{aligned} \quad (4.32)$$

The antisymmetric part of this expression gives the orthogonal splitting of the 2-form $F_{\mu\nu}$ which is

$$F_{\mu\nu} = -Y A_{[\mu} n_{\nu]} + n_{[\mu} D_{\nu]} Y + D_{[\mu} Y_{\nu]} - n_{[\mu} \mathcal{L}_{\vec{n}} Y_{\nu]}. \quad (4.33)$$

Also the symmetric part of (4.32) is the Killing condition $\nabla_{(\mu} \xi_{\nu)} = 0$. Explicitly one has

$$Y K_{\mu\nu} + D_{(\mu} Y_{\nu)} + n_\mu n_\nu (\mathcal{L}_{\vec{n}} Y - A^\rho Y_\rho) + n_{(\mu} (Y A_{\nu)} - 2K_{\nu)\rho} Y^\rho - D_{\nu)} Y - \mathcal{L}_{\vec{n}} Y_{\nu)}) = 0,$$

from which we deduce

$$\mathcal{L}_{\vec{n}} Y_\mu = A_\mu Y - D_\mu Y - 2K_{\mu\rho} Y^\rho, \quad (4.34)$$

which renders the relation (4.33) in the form

$$F_{\mu\nu} = 2n_{[\mu} K_{\nu]\rho} Y^\rho + D_{[\mu} Y_{\nu]} + 2n_{[\mu} D_{\nu]} Y. \quad (4.35)$$

From this expression we may compute the orthogonal splitting of $F_{\mu\nu}^*$ which is

$$F_{\mu\nu}^* = \varepsilon_{\mu\nu\beta} (K_\alpha{}^\beta Y^\alpha + D^\beta Y) + n_{[\mu} \varepsilon_{\nu]\alpha\beta} D^\alpha Y^\beta, \quad (4.36)$$

where equation (2.5) was used along the way. Inserting equations (4.35) and (4.36) in (4.3) and applying (4.28) on the resulting expression we deduce

$$\mathcal{E}_\mu = K_{\mu\rho} Y^\rho + D_\mu Y - \frac{1}{2} \varepsilon_{\mu\nu\rho} D^\nu Y^\rho, \quad (4.37)$$

which is the required relation.

5. The causal propagation of the Mars-Simon tensor

In this section a proof of the following result is provided.

Theorem 4. *The Mars-Simon tensor $\mathcal{S}_{\mu\nu\rho\lambda}$ propagates causally.*

Proof. According to (4.20) it is enough to show that either $\text{Re}(\mathcal{S}_{\mu\nu\rho\lambda})$ or $\text{Im}(\mathcal{S}_{\mu\nu\rho\lambda})$ propagates causally. We choose to work with the former, so let us set $S_{\mu\nu\rho\lambda} \equiv \text{Re}(\mathcal{S}_{\mu\nu\rho\lambda})$ and define $L_{\mu\nu\lambda} \equiv \text{Re}(\mathcal{J}_{\mu\nu\lambda})$. Equation (4.21) entails

$$\nabla_\alpha S^\alpha{}_{\beta\gamma\delta} = L_{\beta\gamma\delta}. \quad (5.1)$$

Hence, $L_{\beta\gamma\delta}$ is the Weyl current of the Weyl candidate $S_{\mu\nu\rho\lambda}$. To study the causal propagation of $S_{\mu\nu\rho\lambda}$ we follow the general procedure explained in section 3.1. We denote by $B_{\mu\nu\rho\lambda}$ the Bel-Robinson tensor constructed from the Weyl candidate $S_{\mu\nu\rho\lambda}$. To prove our result, we need to find a good estimate for the quantity $n^\nu n^\rho n^\lambda \nabla_\mu B_{\nu\rho\lambda}^\mu$ —see equation (3.19). One can write

$$n^\nu n^\rho n^\lambda \nabla_\mu B_{\nu\rho\lambda}^\mu = L^{\rho\nu\lambda} n^\mu S_{\mu\rho\nu\lambda} + 4L^\mu{}_\rho{}^\nu n^\rho n^\lambda n^\alpha S_{\lambda\mu\alpha\nu}, \quad (5.2)$$

where (3.9) was used. In order to work out the right hand side of equation (5.2) we calculate the orthogonal splitting of $S_{\mu\nu\rho\lambda}$ and $L_{\mu\nu\lambda}$ with respect to n^μ . The tensors $S_{\mu\nu\rho\lambda}$ and $L_{\mu\nu\rho}$ are related to $\mathcal{S}_{\mu\nu\rho\lambda}$ and $\mathcal{J}_{\mu\nu\rho}$ by means of the relations

$$S_{\mu\nu\rho\sigma} = \frac{1}{2}(\mathcal{S}_{\mu\nu\rho\sigma} + \overline{\mathcal{S}}_{\mu\nu\rho\sigma}), \quad L_{\mu\nu\rho} \equiv \frac{1}{2}(\mathcal{J}_{\mu\nu\rho} + \overline{\mathcal{J}}_{\mu\nu\rho}), \quad (5.3)$$

and therefore their orthogonal splittings can be calculated once those of $\mathcal{S}_{\mu\nu\rho\sigma}$ and $\mathcal{J}_{\mu\nu\rho}$ are known. The orthogonal splitting of $\mathcal{S}_{\mu\nu\rho\sigma}$ is given in equation (4.25) and the orthogonal splitting of $\mathcal{J}_{\mu\nu\rho}$ can be calculated from equation (4.23) given that we know the orthogonal splitting of all the quantities which appear in the definition of $\mathcal{J}_{\mu\nu\rho}$ —see (A.1) and (A.2)-(A.5) in appendix Appendix A.1 for the precise formulae. Inserting these orthogonal splittings in (5.2) we get

$$\begin{aligned} n^\nu n^\rho n^\lambda \nabla_\mu B_{\nu\rho\lambda}^\mu &= 12 \operatorname{Re} \left[\frac{1}{\sigma - 1} \left(\mathcal{E}^\mu Y_\mu \mathcal{T}^{\rho\lambda} \overline{\mathcal{T}}_{\rho\lambda} - 2\mathcal{E}^\mu Y^\nu \mathcal{T}_{(\mu}{}^\lambda \overline{\mathcal{T}}_{\nu)\lambda} \right) \right] \\ &\quad - 6i \varepsilon_{\mu\nu\rho} \mathcal{T}^{\lambda\nu} \overline{\mathcal{T}}_\lambda{}^\rho Y \left(\frac{\mathcal{E}^\mu}{\sigma - 1} - \frac{\overline{\mathcal{E}}^\mu}{\overline{\sigma} - 1} \right). \end{aligned} \quad (5.4)$$

Now, using the equations (A.13)-(A.14) of appendix Appendix A.2 we can deduce the property

$$B_{\alpha\mu\nu\rho}(\mathcal{E}^\alpha n^\nu n^\rho Y^\mu - \mathcal{E}^\alpha n^\mu n^\nu n^\rho Y) = -i \varepsilon_{\mu\nu\alpha} \mathcal{E}^\mu \mathcal{T}^{\rho\nu} \overline{\mathcal{T}}_\rho{}^\alpha Y + \mathcal{E}^\mu Y_\mu \mathcal{T}^{\alpha\beta} \overline{\mathcal{T}}_{\alpha\beta} - 2\mathcal{E}^\mu Y^\nu \mathcal{T}_{(\nu}{}^\rho \overline{\mathcal{T}}_{\mu)\rho},$$

which when combined with (5.4) yields

$$n^\nu n^\rho n^\sigma \nabla_\mu B_{\nu\rho\sigma}^\mu = \operatorname{Re} \left[\frac{12}{1 - \sigma} B_{\mu\nu\alpha\beta} (\mathcal{E}^\mu n^\beta n^\nu n^\alpha Y - \mathcal{E}^\mu n^\alpha n^\beta Y^\nu) \right]. \quad (5.5)$$

Equation (5.5) is the key to get the required estimate. To obtain it, let us introduce an orthonormal frame $E = \{e_0^\mu, e_1^\mu, e_2^\mu, e_3^\mu\}$ with $n^\mu = e_0^\mu$. Since $\mathcal{E}^\mu, Y^\mu$ are spatial vectors they can be written as a linear combination of the vectors $e_1^\mu, e_2^\mu, e_3^\mu$. Therefore, point (ii) of theorem 2 entails the estimates

$$\operatorname{Re} \left[\frac{12}{1 - \sigma} B_{\mu\nu\alpha\beta} \mathcal{E}^\mu n^\beta n^\nu n^\alpha Y \right] \leq m_1 B_{\mu\nu\alpha\beta} n^\mu n^\nu n^\alpha n^\beta, \quad (5.6a)$$

$$-\operatorname{Re} \left[\frac{12}{1 - \sigma} B_{\mu\nu\alpha\beta} \mathcal{E}^\mu n^\alpha n^\beta Y^\nu \right] \leq m_2 B_{\mu\nu\alpha\beta} n^\mu n^\nu n^\alpha n^\beta, \quad (5.6b)$$

for some scalar continuous functions m_1, m_2 defined on $\mathcal{K}$. Using these estimates in equation (5.5) we obtain our final estimate $n^\nu n^\rho n^\sigma \nabla_\mu B_{\nu\rho\sigma}^\mu \leq m_3 B_{\mu\nu\alpha\beta} n^\mu n^\nu n^\alpha n^\beta$ where

m_3 is the upper bound of $m_1 + m_2$ in the compact set $\mathcal{K}$ and thus (3.19) can be written in the form

$$0 \leq W'(t) \leq (m_3 + 3m_0)W(t), \quad \forall t \in [0, t_1]. \quad (5.7)$$

Since $W(0) = 0$, then an application of Grönwall's lemma enables us to conclude that $W(t) = 0$, $\forall t \in [0, t_1]$ which, according to the considerations made at the end of section 3.1 proves the causal propagation. $\square$

Remark 2. We stress that the fact that $\mathcal{S}_{\mu\nu\rho\lambda}$ satisfies a wave equation and the property (4.21) do not, by themselves, suffice to show the causal propagation of this tensor. If we impose that $\mathcal{S}_{\mu\nu\rho\lambda}$ and some directional derivative of this quantity vanish on a spacelike hypersurface $\mathcal{B}$ then using the wave equation and standard results on hyperbolic partial differential equations allow us to conclude that $\mathcal{S}_{\mu\nu\rho\lambda}$ vanishes on a neighbourhood of $\mathcal{B}$. In doing so, extra conditions needed to be imposed so that $\mathcal{S}_{\mu\nu\rho\lambda}$ is indeed zero in $D(\mathcal{B})$. This approach leads, in principle, to a weaker result than the one given in theorem 4.

Consider now any vacuum spacetime admitting a Killing vector ξ^μ and use ξ^μ to construct the Mars-Simon tensor $\mathcal{S}_{\mu\nu\rho\lambda}$ and as above, define its real part $S_{\mu\nu\rho\lambda}$ and the Bel-Robinson tensor $B_{\mu\nu\rho\lambda}$ of $S_{\mu\nu\rho\lambda}$. For any unit timelike vector n^μ , define the positive scalar

$$\Phi(\vec{n}, \vec{\xi}) \equiv B_{\mu\nu\rho\alpha} n^\mu n^\nu n^\rho n^\alpha. \quad (5.8)$$

One can find an explicit expression for $\Phi(\vec{n}, \vec{\xi})$ if we use the property —see equation (A.12) in Appendix A.2—

$$B_{\mu\nu\rho\alpha} n^\mu n^\nu n^\rho n^\alpha = \mathcal{T}_{\mu\nu} \bar{\mathcal{T}}^{\mu\nu} = 4\mathcal{E}^{\mu\nu} \bar{\mathcal{E}}_{\mu\nu} - 24\text{Re} \left[\frac{\mathcal{E}^\mu \mathcal{E}^\nu \mathcal{E}_{\mu\nu}}{\sigma - 1} \right] - 12 \left| \frac{\mathcal{E}^\mu \mathcal{E}_\mu}{\sigma - 1} \right|^2 + 36 \left| \frac{\mathcal{E}^\mu \bar{\mathcal{E}}_\mu}{\sigma - 1} \right|^2, \quad (5.9)$$

which shows that $\Phi(\vec{n}, \vec{\xi})$ can be expressed in terms of $\mathcal{E}_{\mu\nu}$, $\mathcal{E}_\mu$ and σ exclusively. When $\Phi(\vec{n}, \vec{\xi})$ vanishes on a Cauchy hypersurface, the causal propagation of $\mathcal{S}_{\mu\nu\rho\lambda}$ entails the following result.

Theorem 5. *Let $(\mathcal{M}, g_{\mu\nu})$ be a globally hyperbolic vacuum spacetime and suppose that $\mathcal{M}$ admits a Killing vector ξ^μ . If there are a Cauchy hypersurface $\mathcal{B} \subset \mathcal{M}$ and a real constant M such that $\Phi(\vec{N}, \vec{\xi})|_{\mathcal{B}} = 0$ and $\Xi(M, \vec{\xi})|_{\mathcal{B}} = 0$ where $\vec{N}$ is the unit timelike normal to $\mathcal{B}$, then $\mathcal{M}$ is locally isometric to the Kerr spacetime.*

Proof. If $\Phi(\vec{N}, \vec{\xi})$ is zero on $\mathcal{B}$ then by (5.8) $B_{\mu\nu\rho\alpha} N^\mu N^\nu N^\rho N^\alpha|_{\mathcal{B}} = 0$. Therefore, remark 1 leads us to

$$\mathcal{S}_{\mu\nu\rho\alpha}|_{\mathcal{B}} = 0.$$

Using that the Mars-Simon tensor propagates causally, one deduces that $\mathcal{S}_{\mu\nu\rho\alpha} = 0$ on $\mathcal{M}$. In particular, this implies that (4.13) is rendered in the form

$$\nabla_\mu \Xi(M, \vec{\xi}) = \frac{4\sigma_\mu \Xi(M, \vec{\xi})}{\sigma - 1}, \quad (5.10)$$

which can be integrated to yield $\Xi = A(1 - \sigma)^4$ for some complex constant A . The condition $\Xi(M, \vec{\xi})|_{\mathcal{B}} = 0$ implies that $A = 0$ and hence $\Xi(M, \vec{\xi})$ vanishes on $\mathcal{M}$.

Theorem 3 can now be applied and we conclude that $\mathcal{M}$ is locally isometric to the Kerr spacetime. $\square$

6. Application: characterization of Kerr initial data

We explain in this section how to use previous results to find conditions which ensure that the data development of a vacuum initial data set is isometric to a subset of the Kerr spacetime. The basic idea behind is to express the result of theorem 5 in terms of conditions on a vacuum initial data set (Σ, h_{ij}, K_{ij}) .

Theorem 6 (Kerr initial data). *Let (Σ, h_{ij}, K_{ij}) be a vacuum initial data set and assume that there exist two scalar fields $\tilde{Y}$, $\tilde{\sigma}$, a vector field $\tilde{Y}_j$ and a real constant M , all defined on Σ , fulfilling the following conditions*

$$\mathcal{E}_j \mathcal{E}^j = \frac{1}{4} M^2 (1 - \tilde{\sigma})^4, \quad (6.1a)$$

$$4\mathcal{E}^{ij} \bar{\mathcal{E}}_{ij} - 24 \operatorname{Re} \left[\frac{\mathcal{E}^i \mathcal{E}^j \mathcal{E}_{ij}}{\tilde{\sigma} - 1} \right] - \frac{3}{4} M^4 |1 - \tilde{\sigma}|^6 + 36 \left| \frac{\mathcal{E}^i \bar{\mathcal{E}}_i}{\tilde{\sigma} - 1} \right|^2 = 0, \quad (6.1b)$$

$$D_j \tilde{\sigma} = 2\tilde{Y} K_{jl} \tilde{Y}^l - 2 \, i \, \varepsilon_{jml} (K_k{}^l \tilde{Y}^k \tilde{Y}^m + \tilde{Y}^m D^l \tilde{Y}) + 2\tilde{Y} D_j \tilde{Y} + 2\tilde{Y}^l D_{[l} \tilde{Y}_{j]}, \quad (6.1c)$$

$$D_{(i} \tilde{Y}_{j)} + \tilde{Y} K_{ij} = 0, \quad (6.1d)$$

$$D_i D_j \tilde{Y} + \mathcal{L}_{\tilde{Y}^l} K_{ij} = \tilde{Y} (r_{ij} + K K_{ij} - 2K_{il} K_j{}^l), \quad (6.1e)$$

where

$$\mathcal{E}_j \equiv K_{jk} \tilde{Y}^k + D_j \tilde{Y} - \frac{1}{2} i \, \varepsilon_{jkl} D^k \tilde{Y}^l, \quad (6.2a)$$

$$\mathcal{E}_{jk} \equiv \frac{1}{2} (E_{jk} - i \, B_{jk}). \quad (6.2b)$$

Then the data development $(\mathcal{M}, g_{\mu\nu})$ of (Σ, h_{ij}, K_{ij}) is locally isometric to an open subset of the Kerr spacetime.

Remark 3. The conditions (6.1d) and (6.1e) are the *Killing initial data* (KID) equations. It is well-known that if both a solution $(\tilde{Y}, \tilde{Y}_i)$ and the initial data set (Σ, h_{ij}, K_{ij}) are *suitably* smooth, then the development of the initial data (Σ, h_{ij}, K_{ij}) possesses a Killing vector such that the pull-back of its restriction to Σ coincides with the given Killing data —see for example, [9, 13], and in particular [8] for an explicit set of smoothness conditions on the Killing data. Alternatively, if the Killing data is *transversal*, that is $\tilde{Y} \neq 0$, then one can make use of the concept of Killing development to ensure the existence of a spacetime with a Killing vector associated to the Killing data under question —see [1]. Again, under suitable regularity conditions it can be shown that the Killing development always includes the Cauchy development of (Σ, h_{ij}, K_{ij}) —see again [9]: the Killing development is unique and maximal among the class of spacetimes containing the relevant Cauchy data and a Killing vector with non-closed orbits; hence, if the Cauchy development has a timelike Killing vector, then it must be a subset of the Killing development.

Proof. Let $\mathcal{M}$ be the data development of the vacuum initial data set (Σ, h_{ij}, K_{ij}) and let ξ^μ be the Killing vector mentioned in remark 3. As usual consider a foliation $\{\Sigma_t\}$, $t \in I$, of $\mathcal{M}$ with $\Sigma_0 = \phi(\Sigma)$ and let n^μ be the unit timelike normal to the leaves of this foliation. Let us adopt the formalism and notation explained in section 4 with ξ^μ being the Killing vector. Then the orthogonal splitting of ξ^μ is $\xi^\mu = -Yn^\mu + Y^\mu$. As explained in remark 3 we have the property

$$\phi^*Y = \tilde{Y}, \quad \partial_j \phi^\mu Y_\mu = \tilde{Y}_j. \quad (6.3)$$

which combined with equations (6.2a) and (4.37) entails $\partial_j \phi^\mu(\mathcal{E}_\mu) = \mathcal{E}_j$. The orthogonal splitting of σ_μ is calculated on one hand from equation (4.6) using the expressions (4.27), (4.37) and the property $\xi^\mu = -Yn^\mu + Y^\mu$. The result is

$$\sigma_\mu = s_\mu + sn_\mu, \quad (6.4)$$

where

$$\begin{aligned} s &\equiv -2K_{\mu\nu}Y^\mu Y^\nu - 2Y^\mu D_\mu Y + i \varepsilon_{\mu\rho\nu} Y^\mu D^\rho Y^\nu, \\ s_\mu &\equiv 2YK_{\mu\rho}Y^\rho - 2i \varepsilon_{\mu\nu\rho}(K_\alpha{}^\rho Y^\alpha Y^\nu + Y^\nu D^\rho Y) + 2YD_\mu Y + 2Y^\nu D_{[\nu}Y_{\mu]}. \end{aligned} \quad (6.5)$$

On the other hand, the orthogonal splitting of σ_μ can be calculated from the relation $\sigma_\mu = \nabla_\mu \sigma$ yielding

$$s = -\mathcal{L}_{\tilde{n}}\sigma, \quad s_\mu = D_\mu \sigma. \quad (6.6)$$

Combining the previous equation with the pull-back of equation (6.5) under ϕ and (6.1c) we deduce

$$D_j(\tilde{\sigma}) = D_j(\phi^*(\sigma)). \quad (6.7)$$

Upon a suitable choice of the Ernst potential σ , this equation can be integrated on Σ to give $\phi^*(\sigma) = \tilde{\sigma}$. Next consider the Mars-Simon tensor $\mathcal{S}_{\mu\nu\rho\lambda}$ constructed with the Killing vector ξ^μ and calculate its orthogonal splitting with respect to n^μ —equation (4.25). Using (4.30) and our previous results, we get

$$\begin{aligned} \phi^*(\Phi(\vec{n}, \vec{\xi})) &= \phi^*(\mathcal{T}_{\mu\nu}\overline{\mathcal{T}}^{\mu\nu}) \\ &= \left(2\overline{\mathcal{E}}_{jk} + \frac{2}{1-\tilde{\sigma}}(3\mathcal{E}_j\mathcal{E}_k - h_{jk}\mathcal{E}_l\mathcal{E}^l)\right) \left(2\mathcal{E}^{jk} + \frac{2}{1-\tilde{\sigma}}(3\overline{\mathcal{E}}^j\overline{\mathcal{E}}^k - h^{jk}\overline{\mathcal{E}}_m\overline{\mathcal{E}}^m)\right). \end{aligned} \quad (6.8)$$

Expanding this product and using conditions (6.1a)-(6.1b) we conclude

$$\Phi(\vec{n}, \vec{\xi})|_{\phi(\Sigma)} = 0. \quad (6.9)$$

Moreover, equation (6.1a) implies $\Xi(M, \vec{\xi})|_{\phi(\Sigma)} = 0$ —see (4.31). Thus, theorem 5 applies and therefore we conclude that $(\mathcal{M}, g_{\mu\nu})$ must be locally isometric to an open subset of the Kerr spacetime. $\square$

6.1. Schwarzschild initial data

Theorem 6 asserts that under certain conditions the development of a vacuum initial data set is isometric to an open subset of the Kerr spacetime but nothing is said as to the resulting development being one of the specializations of the Kerr spacetime: the Schwarzschild solution. In this section we show the form which theorem 6 takes in that case. This complements previous work of the present authors about the same issue [16].

To state our result we need a preliminary lemma whose proof can be found in [6].

Lemma 1. *Let (Σ, h_{ij}, K_{ij}) be a vacuum initial data set and assume that there exist a scalar $\tilde{Y}$ and a vector field $\tilde{Y}_j$, all defined on Σ , fulfilling the following conditions*

$$\tilde{Y} D_{[i} \tilde{Y}_{j]} + 2\tilde{Y}_{[i} D_{j]} \tilde{Y} + 2\tilde{Y}_{[i} K_{j]l} \tilde{Y}^l = 0, \quad (6.10a)$$

$$\tilde{Y}_{[i} D_j \tilde{Y}_{k]} = 0. \quad (6.10b)$$

Then there exists an integrable Killing vector ξ^μ in the data development of (Σ, h_{ij}, K_{ij}) .

With the aid of this lemma we can now derive conditions guaranteeing that the development of a vacuum initial data set is isometric to an open subset of the Schwarzschild spacetime

Theorem 7 (Schwarzschild initial data). *Let (Σ, h_{ij}, K_{ij}) be a vacuum initial data set and assume that there exist two scalar fields $\tilde{Y}$, $\tilde{\sigma}$ and a vector field $\tilde{Y}_j$, all defined on Σ , fulfilling the conditions of theorem 6 with (6.1d)-(6.1e) replaced by (6.10a)-(6.10b). Then the data development $\mathcal{M}$ is locally isometric to an open subset of the Schwarzschild spacetime.*

Proof. The first step in the proof of this theorem is to use lemma 1 to show that the data development $\mathcal{M}$ admits an integrable Killing vector ξ^μ . The remaining part of the proof follows the same pattern as that of theorem 6 and therefore we reach the conclusion that $\mathcal{M}$ is isometric to an open subset of a specialization of the Kerr spacetime possessing an integrable Killing vector which, as is well known, is the Schwarzschild spacetime. $\square$

Remark 4. It is worth recalling that the analysis of [16] has provided an explicit formula for a *timelike KID candidate* in terms of concomitants of the initial data (h_{ij}, K_{ij}) . Hence, a combination of theorem 7 with the such timelike KID candidate renders an algorithmic characterization of Schwarzschild initial data alternative to the one given in [16].

7. Conclusions

The main result obtained in this article, theorem 6 provides a characterization of initial data for the Kerr spacetime by first requiring that a set of overdetermined elliptic differential equations—the KID equations—admit a solution; and then in turn requiring that certain objects constructed out of the solutions to the KID equation satisfy very particular relations. This characterization fails to be algorithmic in two ways. First, it is in practise a non-trivial problem to verify whether a certain initial

data set admits a KID or not. A pointwise procedure for doing this has been discussed in [2]. Alternatively, one could resort to a non-time symmetric generalization of the geometric invariant characterizing static time symmetric initial data sets constructed in [10] —which requires solving a fourth-order elliptic partial differential equation. In any case —and this leads to the second way the result fails to be algorithmic— even if one could ascertain the existence of a KID on the initial data, one requires to know the KID in an explicit manner in order to be able to verify the conditions (6.1a), (6.1b) and (6.1c). It is plausible that from an hypothetical characterization of the Kerr spacetime in terms of, say, concomitants of the Weyl tensor —analogous to that for the Schwarzschild spacetime given in [15]— one would be able to construct, following the ideas of [16], KID candidates for the initial data set (Σ, h_{ij}, K_{ij}) . If such a KID candidate were available, then theorem 6 would constitute a powerful tool to analyse the behaviour of dynamical black hole spacetimes, both from an analytic and a numerical point of view.

Acknowledgements

We would like to thank JMM Senovilla, M Mars, W Simon and CM Losert-VK for useful remarks on the manuscript. JAVK is supported by an EPSRC Advanced Research Fellowship. AGP gratefully acknowledges financial support from the Spanish “Ministerio de Ciencia e Innovación” under the postdoctoral grant EX-2006-0092.

Appendix A. Technical details about the calculations

Appendix A.1. The orthogonal splitting of $\mathcal{J}_{\mu\nu\rho}$

Given the symmetries of the Weyl current $\mathcal{J}_{\mu\nu\rho}$ it is immediate to realise that its orthogonal splitting must take the form

$$\mathcal{J}_{\mu\nu\rho} = n_\mu \mathcal{Z}_{\nu\rho} + n_{[\rho} \mathcal{X}_{\nu]\mu} + n_\mu n_{[\nu} \mathcal{V}_{\rho]} + \mathcal{P}_{\mu\nu\rho}. \quad (\text{A.1})$$

It takes a greater effort, however, to find the explicit expression of the spatial tensors $\mathcal{Z}_{\mu\nu}$, $\mathcal{X}_{\mu\nu}$, $\mathcal{V}_\mu$, $\mathcal{P}_{\mu\nu\rho}$. These expressions are found by inserting into equation (4.23) the orthogonal splitting of $\mathcal{S}_{\mu\nu\rho\lambda}$ shown in (4.25), the orthogonal splitting of $\mathcal{F}_{\mu\nu}$ shown in (4.27) and the relation $\xi^\mu = -Yn^\mu + Y^\mu$. A computer calculation yields the results

$$\mathcal{Z}_{\mu\nu} \equiv \frac{4}{1-\sigma} (\text{i} Y \mathcal{E}^\alpha \mathcal{T}_{[\mu}{}^\rho \varepsilon_{\nu]\alpha\rho} + 3 \mathcal{E}_{[\mu} \mathcal{T}_{\nu]\rho} Y^\rho + Y_{[\nu} \mathcal{T}_{\mu]\alpha} \mathcal{E}^\alpha - \text{i} \varepsilon_{\mu\nu\rho} \mathcal{E}^\alpha \mathcal{T}_\alpha{}^\rho Y), \quad (\text{A.2})$$

$$\begin{aligned} \mathcal{X}_{\mu\nu} \equiv \frac{1}{1-\sigma} \Big(& \mathcal{E}^\alpha (4\text{i} Y (\mathcal{T}_\alpha{}^\rho \varepsilon_{\nu]\mu\rho} - \varepsilon_{\nu\alpha\rho} \mathcal{T}_\mu{}^\rho) + 6 \mathcal{T}_{\nu\mu} Y_\alpha - 2 \mathcal{T}_{\mu\alpha} Y_\nu - 4 \mathcal{T}_{\nu\alpha} Y_\mu + 4 h_{\nu\mu} \mathcal{T}_{\alpha\rho} Y^\rho \\ & + 6 \mathcal{E}_\nu \mathcal{T}_{\mu\alpha} Y^\alpha \Big), \end{aligned} \quad (\text{A.3})$$

$$\mathcal{V}_\mu \equiv \frac{\mathcal{E}^\alpha}{1-\sigma} \left(4\text{i} Y^\rho (\varepsilon_{\mu\lambda(\rho} \mathcal{T}_{\alpha)}{}^\lambda + \varepsilon_{\alpha\rho\lambda} \mathcal{T}_\mu{}^\lambda) - 2 Y \mathcal{T}_{\mu\alpha} \right), \quad (\text{A.4})$$

$$\mathcal{P}_{\mu\nu\rho} \equiv \frac{2}{1-\sigma} \left(2 \text{i} \varepsilon_{\nu\rho\lambda} \mathcal{E}_\mu \mathcal{T}_\alpha{}^\lambda Y^\alpha + 2 \mathcal{E}_{[\rho} (3 Y \mathcal{T}_{\nu]\mu} + \text{i} \varepsilon_{\nu]\mu\lambda} \mathcal{T}_\alpha{}^\lambda Y^\alpha \right)$$

$$\begin{aligned}
& + \mathcal{E}^\alpha \left(-2i \left(\varepsilon_{\rho\alpha\lambda} \mathcal{T}_{[\nu}{}^\lambda Y_{\mu]} + \varepsilon_{\nu\alpha\lambda} \mathcal{T}_{[\mu}{}^\lambda Y_{\rho]} + 2\varepsilon_{\mu\alpha\lambda} \mathcal{T}_{[\nu}{}^\lambda Y_{\rho]} \right. \right. \\
& \left. \left. + Y^\lambda (\varepsilon_{\nu\rho\alpha} \mathcal{T}_{\mu\lambda} + \varepsilon_{\mu\alpha[\nu} \mathcal{T}_{\rho]\lambda}) \right) \right. \\
& \left. + 2h_{\mu[\nu} \left(\mathcal{T}_{\rho]\alpha} Y + i Y^\lambda (\varepsilon_{\rho]\lambda\delta} \mathcal{T}_\alpha{}^\delta + \mathcal{T}_{\rho]}{}^\delta \varepsilon_{\alpha\lambda\delta} + \varepsilon_{\rho]\alpha\delta} \mathcal{T}_\lambda{}^\delta) \right) \right). \quad (\text{A.5})
\end{aligned}$$

Appendix A.2. The orthogonal splitting of $B_{\mu\nu\rho\sigma}$

In [14] the orthogonal splitting of the Bel-Robinson tensor was calculated and the different parts resulting from such splitting were studied in detail. The calculations are similar for the Bel-Robinson tensor of a Weyl candidate and therefore we only need to adapt these results for the particular case of $S_{\mu\nu\rho\lambda}$. In this case the orthogonal splitting of $B_{\mu\nu\rho\lambda}$ reads —see section (5.1) of [14]—

$$B_{\mu\nu\rho\lambda} = W n_\mu n_\nu n_\rho n_\lambda + 4P_{(\mu} n_\nu n_\rho n_{\lambda)} + 6t_{(\mu\nu} n_\rho n_{\lambda)} + 4Q_{(\mu\nu\rho} n_{\lambda)} + t_{\mu\nu\rho\lambda}, \quad (\text{A.6})$$

where the different parts of this splitting are given by

$$W \equiv e_{\alpha\beta} e^{\alpha\beta} + b_{\alpha\beta} b^{\alpha\beta}, \quad (\text{A.7})$$

$$P_\mu \equiv 2b_\alpha{}^\lambda e_{\rho\lambda} \varepsilon_\mu{}^{\alpha\rho}, \quad (\text{A.8})$$

$$t_{\mu\nu} \equiv W h_{\mu\nu} - 2(b_\mu{}^\rho b_{\nu\rho} + e_\mu{}^\rho e_{\nu\rho}), \quad (\text{A.9})$$

$$Q_{\mu\nu\rho} \equiv h_{\mu\nu} P_\rho - 2(b_{\nu\alpha} e_{\mu\beta} + b_{\mu\alpha} e_{\nu\beta}) \varepsilon_\rho{}^{\alpha\beta}, \quad (\text{A.10})$$

$$\begin{aligned}
t_{\mu\nu\rho\lambda} \equiv & 4(b_{\mu\nu} b_{\rho\lambda} + e_{\mu\nu} e_{\rho\lambda}) - h_{\rho\lambda} t_{\mu\nu} + 2h_{\nu(\lambda} t_{\rho)\mu} + 2h_{\mu(\lambda} t_{\rho)\nu} - h_{\mu\nu} t_{\rho\lambda} + \\
& + W(h_{\mu\nu} h_{\rho\lambda} - 2h_{\mu(\rho} h_{\lambda)\nu}), \quad (\text{A.11})
\end{aligned}$$

where $e_{\mu\nu}$ and $b_{\mu\nu}$ are respectively the *electric* and *magnetic* parts of the Weyl candidate $S_{\mu\nu\lambda\rho}$. From (A.6) and (A.10) we get

$$B_{\mu\nu\rho\lambda} n^\mu n^\nu n^\rho n^\lambda = \mathcal{T}_{\rho\lambda} \overline{\mathcal{T}}^{\rho\lambda} \quad (\text{A.12})$$

$$B_{\mu\nu\rho\lambda} n^\nu n^\rho n^\lambda = -n_\mu \mathcal{T}^{\rho\lambda} \overline{\mathcal{T}}_{\rho\lambda} + i \varepsilon_{\mu\nu\rho} \mathcal{T}^{\lambda\nu} \overline{\mathcal{T}}_\lambda{}^\rho. \quad (\text{A.13})$$

$$B_{\mu\nu\rho\lambda} n^\rho n^\lambda = -2\mathcal{T}_{(\mu}{}^\rho \mathcal{T}_{\nu)\rho} + (n_\mu n_\nu + h_{\mu\nu}) \mathcal{T}^{\rho\lambda} \overline{\mathcal{T}}_{\rho\lambda} - 2i \mathcal{T}^{\alpha\rho} \overline{\mathcal{T}}_\alpha{}^\lambda \varepsilon_{\rho\lambda(\mu} n_{\nu)}, \quad (\text{A.14})$$

where the relations

$$e_{\mu\nu} = \mathcal{T}_{\mu\nu} + \overline{\mathcal{T}}_{\mu\nu}, \quad b_{\mu\nu} = i (\mathcal{T}_{\mu\nu} - \overline{\mathcal{T}}_{\mu\nu}),$$

need to be used.

References

- [1] R. Beig and P. T. Chruściel, *Killing initial data*, Class. Quantum Grav. **14**, A83 (1997).
- [2] R. Beig and P. T. Chruściel, *KIDs are non-generic*, Ann. H. Poincaré **6**, 155 (2005).
- [3] L. Bel *Sur la radiation gravitationnelle*, C. R. Acad. Sci. Paris **247**, 1094 (1958).
- [4] G. Bergqvist and J. M. M. Senovilla, *On the causal propagation of fields*, Class. Quantum Grav. **16**, L55 (1999).
- [5] M. A. G. Bonilla and J.M.M. Senovilla, *A very simple proof of the causal propagation of gravity in vacuum*, Phys. Rev. Lett. **78**, 783 (1997).
- [6] A. Carrasco and M. Mars, *On marginally outer trapped surfaces in stationary and static spaces*, Class. Quantum Grav. **25**, 055011 (2008).

- [7] P. T. Chruściel, *Uniqueness of stationary, electro-vacuum black holes revisited* <http://www.arxiv.org/abs/gr-qc/9610010>, (1996).
- [8] P. T. Chruściel, *On the uniqueness in the large of solutions of Einstein's equations ("Strong Cosmic Censorship")*, Centre for Mathematics and its applications, Australian National University, 1991.
- [9] B. Coll, *On the evolution equations for Killing fields*, J. Math. Phys. **18**, 1918 (1977).
- [10] S. Dain, *A new geometric invariant on initial data for the Einstein equations*, Phys. Rev. Lett. **93**, 101101 (2006).
- [11] G. F. R. Ellis, *Cargèse lectures in Physics*, vol 6 (New York: Gordon and Breach) page 1 (1973).
- [12] E.ourgoulhon, *3 + 1 formalism and bases of numerical relativity*, <http://www.arxiv.org/abs/gr-qc/0703035>, (2007).
- [13] V. Moncrief, *Spacetime symmetries and linearization stability of the Einstein equations. I.*, J. Math. Phys. **16**, 493 (1975).
- [14] A. García-Parrado, *Dynamical laws of superenergy in General Relativity*, Class. Quantum Grav. **25**, 015006 (2008).
- [15] J. J. Ferrando and J. A. Sáez, *An intrinsic characterization of the Schwarzschild metric*, Class. Quantum Grav. **15**, 1323 (1998).
- [16] A. García-Parrado and J.A. Valiente Kroon, *Initial data sets for the Schwarzschild spacetime*, Phys. Rev. D **75**, 024027 (2007).
- [17] A. García-Parrado and J.A. Valiente Kroon, *Killing spinor initial data sets* J. Geom. Phys. doi.10.1016/j.geomphys.2008.04.003 (2008).
- [18] M. Heusler, *Black Hole Uniqueness Theorems*, Cambridge University Press, Cambridge, (1996).
- [19] A. D. Ionescu and S. Klainerman, *On the uniqueness of smooth, stationary black holes in vacuum*, <http://arxiv.org/abs/0711.0040>, (2007).
- [20] S. Klainerman and F. Nicolò, *On local and global aspects of the Cauchy problem in general relativity*, Class. Quantum Grav. **16**, R73-R157, (1999).
- [21] M. Mars, *A spacetime characterization of the Kerr metric*, Class. Quantum Grav. **16**, 2507 (1999).
- [22] M. Mars, *Uniqueness properties of the Kerr Metric*, Class. Quantum Grav. **17**, 3353 (2000).
- [23] J.M. Martín-García see <http://metric.iem.csic.es/Martin-Garcia/xAct>
- [24] J.M. Martín-García *xPerm: fast index canonicalization for tensor computer algebra* <http://www.arxiv.org/abs/0803.0862>, (2008)
- [25] J. M. M. Senovilla, *Superenergy tensors*, Class. Quantum Grav. **17**, 2799 (2000).
- [26] J. M. M. Senovilla, *Symmetric hyperbolic systems for a large class of fields in arbitrary dimension*, Gen. Rel. Grav. **39**, 361 (2007).
- [27] W. Simon, *Characterizations of the Kerr metric*, Gen. Rel. Grav. **16**, 465 (1984).
- [28] J. A. Valiente Kroon, *A characterisation of Schwarzschild initial data*, Phys. rev. D **72**, 084003 (2005).
- [29] R. Wald, *General Relativity*, The University of Chicago, Chicago (1984).